

ORIGINAL RESEARCH

Maternal immune activation alters adult behavior, intestinal integrity, gut microbiota and the gut inflammation

Wenqiang Li^{1,2,3}  | Mengxue Chen^{1,2,3}  | Xia Feng^{1,2,3} | Meng Song^{1,2,3} |
Minglong Shao^{1,2,3} | Yongfeng Yang^{1,2,3} | Luwen Zhang^{1,2,3} | Qing Liu^{1,2,3} |
Luxian Lv^{1,2,3,4} | Xi Su^{1,2,3}

¹Henan Mental Hospital, The Second Affiliated Hospital of Xinxiang Medical University, Xinxiang, China

²Henan Key Lab of Biological Psychiatry of Xinxiang Medical University, Xinxiang, China

³International Joint Research Laboratory for Psychiatry and Neuroscience of Henan, Xinxiang, China

⁴Henan Province People's Hospital, Zhengzhou, China

Correspondence

Xi Su and Luxian Lv, Department of Psychiatry of the Second Affiliated Hospital of Xinxiang Medical University, No.388, Jianshe Middle Road, 453002 Xinxiang, China.
Emails: suxi198919@163.com; lvx928@126.com

Funding information

National Natural Science Foundation of China, Grant/Award Number: 81972152, 82001407 and U1904130; High Scientific and Technological Research Fund of Xinxiang Medical University, Grant/Award Number: 2017ZDCG-04; Medical Science and Technology Research Project of Henan Province, Grant/Award Number: 2018020371, LHGJ20190476 and LHGJ20190482; The Open Fund of Henan Key lab of Biological Psychiatry, Grant/Award Number: ZDSYS2018003 and ZDSYS2018008; Key Scientific Research Project of Henan Higher Education Institutions, Grant/Award Number: 20B320016

Abstract

Background: Schizophrenia is characterized by several core behavioral features, in which the gastrointestinal symptoms are frequently reported. Maternal immune activation (MIA) has been developed in a rodent model to study neurodevelopmental disorders such as schizophrenia. However, the changes in the gut environment of MIA rats remain largely unknown.

Methods: 10 mg/kg of polyinosinic:polycytidylic acid (Poly I:C) on gestational day 9 was intravenously administered to rats to induce MIA in order to assess changes in behavior, the intestinal barrier and microbiota in offspring.

Results: Maternal immune activation offspring shown increased anxiety as indicated by reduced exploration of central area in open field test and decreased exploration of open arms in elevated plus test. Cognitive impairment of MIA offspring was confirmed by reduced exploration of novel arm in Y maze test and deficiency of PPI. Intestinal muscle thickness became thinner and some specific microbial anomalies previously identified clinically were observed in MIA offspring. In addition, an increase of inflammatory responses was found in the gut of MIA offspring.

Conclusions: Maternal immune activation alters behavior, intestinal integrity, gut microbiota and the gut inflammation in adult offspring.

KEYWORDS

behavior, gut, inflammation, maternal immune activation, microbiota

Wenqiang Li and Mengxue Chen should be considered joint first author.

Luxian Lv and Xi Su should be considered joint senior author.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Brain and Behavior* published by Wiley Periodicals LLC

1 | BACKGROUND

Schizophrenia is a common and severe neurodevelopmental disease that affects approximately 1% of the population. This disease is manifested by a disruption in cognition and emotion, along with other negative and positive symptoms. Although schizophrenia is characterized by core behavioral impairment, gastrointestinal (GI) symptoms are frequently reported (Nemani et al., 2015). A proportion of schizophrenic patients exhibit dysbiosis of the intestinal microbiota, with some demonstrate an increase in intestinal permeability (Akhondzadeh, 2019; Golofast & Vales, 2020; Nguyen et al., 2019).

The maternal immune activation (MIA) model of schizophrenia has been established based on several epidemiological studies (Brown et al., 2004; Mednick et al., 1988; Meyer & Feldon, 2010). In addition to the changes in neuromorphology, electrophysiology, neurochemistry and neuronal structure, the adult offspring of MIA rodent often exhibit behavioral alterations and other characteristics of schizophrenia (Meyer, 2014; Meyer et al., 2008). One of the most common technique for MIA induction is the supplementation of double-stranded RNA analogue to the pregnant rodents (Meyer, 2014). Several reports have demonstrated that the offspring of pregnant dams exposed to polyinosinic:polycytidylic acid (poly I:C) display schizophrenia-like behavior, including cognitive deficits, as well as exhibit neuroimmunological and brain morphological abnormalities (Bergdolt & Dunaevsky, 2019; Meyer, 2014; Murray et al., 2019). Hence, the poly I:C-induced MIA model has emerged as a valuable research tool for the assessment of schizophrenia.

Commensal bacteria may induce various behavioral features such as emotional, social, and anxiety-like behavior, thereby affecting the neuronal development and function of both mice (Cryan & Dinan, 2012) and humans (Tillisch et al., 2013). Gut microbiota are highly sensitive to early developmental environments, including MIA (Hsiao et al., 2013). More recently, it has been hypothesized intestinal microbiota can affect the host's brain development (Hsiao et al., 2013).

Schizophrenia is a complex mental disorder with unknown etiology. MIA is one of the most ideal models for studying neurodevelopment disorders (Brown & Meyer, 2018). Based on the recent emerging development of a gut-microbiota-brain axis, it is hypothesized that the behavioral features of schizophrenia may be related to GI deficits (Golofast & Vales, 2020). If analogous GI dysfunction exist in MIA rat model, it will be suitable to investigate the pathophysiology of the gut-brain interaction of neuropsychiatry, and potentially suggesting the GI as a new target for psychotherapeutic selection. The aim of this study was to investigate whether MIA causes GI deficiency and schizophrenia-like behaviors. Further investigations are needed to address questions related to the cause and relationship between the gut and the brain: do changes in the GI underlie the pathophysiology of schizophrenia or are they a result thereof?

2 | METHODS

2.1 | Animals

Sprague-Dawley rats (8 weeks old) were supplied by Beijing Vital Rival. The rats were allowed to mate at 3 months old, and the occurrence of mating was verified by copulatory plugs. All rats were housed in ventilated cages at $22 \pm 2^\circ\text{C}$ and $50 \pm 10\%$ humidity, under a normal day/night cycle (lights on from 08:00 to 20:00). All animal handling and experimental procedures were conducted according to the National Institute of Health Guide for the Care and Use of Laboratory Animal. All animal sample collection protocols complied with the current laws of China. All animal procedures performed in this research were in accordance with the Laboratory Animal Ethics Committee of Henan Mental Hospital. (permit number: HMH. No20190806AEC004).

2.2 | MIA during pregnancy

A timed-mating procedure was employed for the construction of MIA model during rat pregnancy. On the 9th day after copulation, the mating dams were randomly assigned into two different groups ($n = 8$ for poly I:C administration group and $n = 8$ for saline administration group). Random numbers were generated using the standard = RAND() function in Microsoft Excel. The rat dams were intravenously injected with 10 mg/kg poly I:C (Sigma-Aldrich) in saline, or an equal amount of saline solution according to previous study (Murray et al., 2019). Three hours after treatment, three rat dams of each group were randomly selected and anesthetized with 50 mg/kg sodium thiopental via intraperitoneal injection. Plasma samples were collected by cardiac puncture using sterile syringes from each rat for measuring the levels of interleukin (IL)-1 β , IL-6 and tumor necrosis factor- α (TNF- α) through the use of enzyme-linked immunosorbent assay (ELISA; R&D Systems).

2.3 | Behavioral testing

Weaning of both pup groups was conducted on postnatal day 21. Previous study suggested sex-specific effects, and particularly male vulnerability, following MIA (Hui et al., 2018; Meehan et al., 2017; Rahman et al., 2017), so male dams were selected for research and no more than two male rats from per litter are used in each group. Ten male rats from each group were randomly selected on postnatal day 60 for a typical behavioral assessment. The open-field, elevated plus-maze, and Y-maze tests were performed with Spain pan Lab Smart version 3.0 software (RWD Life Science.) using an overhead video camera system to automate behavioral testing and provide unbiased data analyses. The open-field and elevated plus-maze tests were adopted to determine anxiety levels and locomotor activities, while Y-maze test was

used to evaluate spatial recognition memory. Then, sensorimotor gating was measured by the prepulse inhibition (PPI) analysis. All these behavioral tests were carried out between 08:00 and 18:00 hr. After each assessment, the apparatus were cleaned with 75% alcohol. All behavioral tests were performed as previously described (Hao et al., 2019).

2.4 | Tissue collection

At adulthood, male rats were anesthetized with 50 mg/kg sodium thiopental via intraperitoneal injection 2 days after behavior testing (behavior testing began at postnatal day 60). Prefrontal cortex and colon tissues from each rat were subjected to mRNA and protein expression analyses. Rats were perfused with saline followed by 4% paraformaldehyde in PBS for hematoxylin and eosin, immunohistochemical and immunofluorescence staining. Stool samples were collected from male adult offspring prior to the start of any behavioral test. The samples were collected in the morning after transferring the animals to freshly disinfected cages.

2.5 | RNA extraction and quantification

Total RNA was extracted with TRIzol (Invitrogen) by following the manufacturer's instructions. Quantitative real-time polymerase chain reaction (qRT-PCR) was conducted on an ABI StepOne RT-PCR System (Applied Biosystems) using a QuantiTect SYBR Green RT-PCR kit (Qiagen, 204245). GAPDH expression was stable between different MIA groups without difference and was chosen as the reference gene of MIA model (Hao et al., 2019). The copy numbers of target mRNAs were normalized to GAPDH mRNA levels in each sample. The primers for RT-PCR are listed in the (Table S1).

2.6 | Protein isolation and western blotting

Protein samples were isolated using radioimmunoprecipitation assay (RIPA) lysis buffer, followed by centrifugation at 12,000 \times g for 15 min at 4°C. After supernatant collection, the protein concentrations were evaluated by the Bio-Rad Coomassie Blue protein assay (Bio-Rad, Hemel Hempstead). Equal amounts of protein (40 μ g) were separated on SDS-PAGE, followed by Western blotting as previously described (Hao et al., 2019).

2.7 | Antibodies and reagents

The primary antibodies used were as follows: anti-IL-1 β (1:1,000; 16806-1-AP; Proteintech), anti-TNF- α (1:1,000; 17590-1-AP; Proteintech), and anti-IL-6 (1:1,000; 21865-1-AP; Proteintech). The secondary horseradish peroxidase (HRP)-conjugated antibodies used were as follows: goat anti-rabbit HRP (1:5,000; 12-348) and goat

anti-mouse (1:5000; sc-2005; Santa Cruz, Insight Biotechnology). Meanwhile, the immunofluorescent CY3 dye-conjugated secondary antibody (1:300; Bost Biotech) was also used.

2.8 | DNA extraction and analysis

Total DNA was isolated from fecal samples using a Fast DNA Stool Mini Kit (QIAGEN). Quantification of DNA yield was performed using a Nanodrop 2000 spectrophotometer. The amplification of six bacteria species (*Bacteroides* spp., *Bifidobacterium* spp., *Clostridium* spp., *Escherichia coli*, *Fusobacterium prausnitzii* and *Lactobacillus* spp.) was conducted on an ABI StepOne RT-PCR System (Applied Biosystems) using a QuantiTect SYBR Green RT-PCR kit (Qiagen, 204245). Bacterial copy numbers were calculated according to the obtained cycle thresholds (CT). The primers for Real-Time PCR detection are listed in the (Table S1).

2.9 | Statistical analysis

The data were analyzed using SPSS Statistics version 20.0 (SPSS Inc.) and were expressed as mean \pm standard error of the mean (SEM). PPI dataset was analyzed using two-way ANOVA (treatment and trials) with repeated measures. Independent student's *t*-tests (two-tailed) were used for other data analysis. A *p*-value of <.05 was considered statistically significance. All experiments were performed at least three times and the representative data were presented.

3 | RESULTS

3.1 | Levels of cytokines in MIA rat model

After administrating 10 mg/kg poly I:C to the pregnant rats, the inflammatory responses were assessed 3 hr following injection. Figure 1 demonstrates the levels of IL-6 ($t = 8.634$, $df=4$, $p = .0010$), IL-1 β ($t = 6.312$, $df=4$, $p = .0032$), and TNF- α ($t = 5.046$, $df=4$, $p = .0073$) in pregnant rats were upregulated in MIA-induced rats compared to saline controls, confirming the activation of maternal immunity by poly I:C.

3.2 | MIA leads to exacerbated anxiety in offspring

Open-field and elevated plus-maze tests were conducted to evaluate the anxiety status of MIA offspring. During the exposure to an open-field arena for 30 min, MIA offspring exhibited a decrease in distance moved within the central area ($t = 2.419$, $df=18$, $p = .0264$), although no differences were observed in distances moved within the whole field (Figure 2a,b). In addition, the animal entry into central areas ($t = 2.625$, $df=18$, $p = .0172$) and time spent in central areas ($t = 2.795$, $df=18$, $p = .0120$) were significantly decreased in




FIGURE 1 Inflammatory cytokine levels in pregnant rats exposed to Poly I:C or saline. The bar graph shows maternal IL-6 (a), IL-1 β (b), and TNF- α (c) blood levels measured 3 hr post 10 mg/kg Poly I:C or saline administration. Independent Student's t-test (two-tailed) was performed for statistical analysis. Results are shown as the mean \pm SEM ($n = 3$ per group). * $p < .05$, ** $p < .01$. The detection was repeated independently for three times

MIA group compared to saline control group (Figure 2c,d). For the elevated plus-maze test, prenatal poly I:C exposure reduced the percentage of time spent in the open arms ($t = 5.330$, $df=18$, $p < .0001$), with fewer entries into the open arms ($t = 3.605$, $df=18$, $p = .0022$) (Figure 2e,f). These findings reveal that MIA offspring display certain anxiety-related behaviors.

3.3 | MIA leads to cognitive impairment in offspring

To examine memory impairment, Y-maze assessment was carried out. Over the course of a 5-min period, poly I:C treatment groups showed fewer entries into the novel arm ($t = 2.936$, $df=18$, $p = .0088$) and decreased the percentage of time spent in the novel arm ($t = 2.190$, $df=18$, $p = .0448$), when compared with saline treated groups (Figure 3a,b).

Further, auditory sensory gating deficit was assessed by PPI test. Compared to control group, the percentage PPI at the prepulse intensities 75, 80, and 85 dB was markedly reduced in MIA rat offspring (Figure 3c).

Altogether, these behavioral findings indicate that MIA can induce schizophrenia-like behavior in rat offspring.

3.4 | MIA induces intestinal impairment in offspring


Subsets of schizophrenia patients are found to exhibit GI abnormalities, including leaky gut syndrome. Therefore, we further determine whether MIA can alter the pathophysiology of the intestines. It was found that the adult MIA offspring with schizophrenia changes displayed significant colon deficits in gut barrier integrity. MIA offspring exhibited hallmark distortions of gut morphology ($t = 5.623$,

$df=58$, $p < .0001$), as indicated by the histological examination of HE-stained sections (Figure 4a). Moreover, severe cellular injury was also observed in the colons of adult MIA offspring, as revealed by TUNEL staining ($t = 2.804$, $df=4$, $p = .0486$), an indicative of marked DNA damage (Figure 4b). Consistent with the defects in intestinal barrier integrity, the levels of the tight junction markers (i.e., ZO-1 and CLDN) were reduced in the colon tissue of adult MIA offspring, as indicated by RT-PCR (claudin: $t = 3.190$, $df=10$, $p = .0110$; zo-1: $t = 3.293$, $df=10$, $p = .0081$)(Figure 4c) and immunohistochemical staining (zo-1: $t = 3.277$, $df=4$, $p = .0306$)(Figure 4d). These observations suggest that intestinal barrier integrity is impaired in these animals.

3.5 | MIA alters gut microbiota composition

Intestinal bacteria play important roles in maintaining intestinal homeostasis and function. Abnormal features associated with dysbiotic microbiome, such as altered bacterial community composition and content, have been observed in schizophrenia patients. To assess whether MIA can induce microbiota alteration, we examined several microorganisms that have been differentially reported in patients with schizophrenia. As illustrated in Figure 5a, the total number bacteria was significantly reduced in MIA offspring, as assessed by 16s rRNA amplification ($t = 2.882$, $df=10$, $p = .0163$). Moreover, the offspring exposed to MIA displayed significantly higher levels of *E. coli* ($t = 2.279$, $df=10$, $p = .0459$; Figure 5b), *Lactobacillus* spp. ($t = 2.534$, $df=10$, $p = .0420$; Figure 5c), *Bifidobacterium* spp. ($t = 2.707$, $df=10$, $p = .0220$; Figure 5d), and *Bacteroides* spp. ($t = 2.406$, $df=10$, $p = .0395$; Figure 5e) when compared with control animals. No significant changes were observed for the fecal *Clostridium coccoides* group (Figure 5f) and *Fusobacterium prausnitzii* spp. (Figure 5g) in MIA

FIGURE 2 Prenatal immune activation induces anxiety behaviors in offspring. (a–d) The open field test. Total distance traveled in the open field arena (a), distance moved in the center arena (b), entries into the center arena (c) and time spent in the center arena (d). (e and f) The elevated plus maze test. Entries into the open arms (e) and time spent in open arms (f); $n = 10$ for each group. Independent Student's *t*-test (two-tailed) was performed for statistical analysis. Data are shown as mean \pm SEM. * $p < .05$, ** $p < .01$



offspring when compared with control animals. These results indicate that common gut microbiota are remarkably altered in MIA offspring.

3.6 | MIA induces intestinal inflammation in MIA offspring

Intestinal permeability is often associated with the changes in immune responses. Notably, the colon tissue of adult MIA offspring displayed elevated mRNA (Figure 6a) and protein expression (Figure 6b,c) levels for IL-1 β and TNF- α , while no change was

observed for IL-6 expression. Immune responses in the prefrontal cortex of MIA offspring were also detected (Figure 6d,e). The changes in intestinal cytokines were not accompanied with immune activation in the prefrontal cortex, as assessed by cytokine detection in the prefrontal cortex (Figure 6d). Through the use of microglia marker Iba1, no significant difference in prefrontal cortex staining was observed between MIA offspring and control animals (Figure 6e). This observation suggests the absence of inflammation in the prefrontal cortex of MIA offspring.

Collectively, it was observed that the adult offspring of MIA dams exhibited increased intestinal permeability and abnormally




FIGURE 3 Prenatal immune activation induces cognitive impairment in offspring. (a and b) The Y-maze test. Entries into the novel arm (a) and time spent in the novel arm (b) were analyzed. (c) % PPI performance at 75, 80, and 85 dB; $n = 10$ for each group. Independent Student's *t* test (two-tailed) was performed for statistical analysis in (a and b); PPI dataset (c) was analyzed using Two-way ANOVA (treatment and trials) with repeated measures. Data are mean \pm SEM. * $p < .05$, ** $p < .01$

expressed levels of cytokines, which are consistent with the characteristics of human schizophrenia.

4 | DISCUSSION

Although the underlying etiology remains unclear, there is substantial evidence of an association between psychiatric disorders and GI disease (Stefano et al., 2018; Van Oudenhove et al., 2010). Animal models are ideal tools for elucidating the mechanisms of human diseases. This study aimed to determine whether the well-characterized MIA model can produce an intestinal environment imbalance concomitant with behavioral abnormalities. Our results indicated that the MIA rat model not only displays well-documented behavioral abnormalities, but also exhibits intestinal integrity disruption, microbial imbalances, and gut inflammation. These data suggest that this model may be a useful tool for elucidating the biological mechanisms associated with GI disorders in psychiatric patients.

Accumulating evidence suggests that exposure to MIA during pregnancy increases the risk of neuropsychiatric diseases, including autism spectrum disorder and schizophrenia (Minakova & Warner, 2018). Epidemiological findings suggest that maternal infection during early-to-mid human pregnancy is more likely to be

associated with long-term developmental brain and behavioral abnormality in the offspring (Brown et al., 2004; Mednick et al., 1988). Different time periods of poly (I:C) exposure can lead to different emergent behaviors and neuropsychiatric changes throughout the offspring life, and infection and related inflammation in the second trimester of gestation can result in long-lasting consequences with regard to brain development and behavior (Meyer, Nyffeler, et al., 2006). However, few reports have suggested that prenatal immune activation during early pregnancy can induce neuropsychiatric disease (Hao et al., 2019; Meyer et al., 2006; Rahman et al., 2017). Previous *in vivo* findings have indicated that prenatal immune activation during early pregnancy, especially on gestation day 9, can induce abnormal behavioral features, including working memory impairment and PPI deficit (Hao et al., 2019). Our results confirmed that early immune activation led to schizophrenia-like behavioral abnormalities in rats. Although psychiatric disorders are typically characterized by core behavioral impairment, GI abnormalities are frequently reported (Sharon et al., 2016). A growing body of literature has linked neurobehavioral symptoms to gut-brain axis dysfunction (Hsiao et al., 2013; Mayer, 2011; Mayer et al., 2014). Subsets of patients exhibit dysbiosis of the gut microbiota, and some demonstrate increased intestinal permeability (Sharon et al., 2016). Recently, several elements of autism spectrum disorder-associated




FIGURE 4 The effects of MIA on the intestinal epithelial barrier in offspring. (a) Representative images of HE staining of colon from MIA offspring were shown; Muscle thickness was analyzed from ten different regions of each sample ($n = 3$ for each group). MT: muscle thickness. (b) Representative images of TUNEL staining of colon from MIA offspring were shown and the percentage of TUNEL positive cell was calculated ($n = 3$ for each group). (c) The mRNA expression of the tight junction markers, claudin and ZO-1 were detected in colon from MIA and control offspring ($n = 6$ for each group). (d) Representative images of IHC staining of claudin and ZO-1 from MIA and control colon offspring were presented, and positive area of immunostain was calculated ($n = 3$ for each group). Independent Student's *t*-test (two-tailed) was performed for statistical analysis. All data are expressed as mean \pm SEM. * $p < .05$, ** $p < .01$. All experiments were repeated independently for three times

and schizophrenia-associated gut-brain axis dysfunction were reproduced in the MIA poly (I:C) model of C57BL6/J mice (Hsiao et al., 2013). MIA and maternal separation could trigger intestinal abnormalities such as increased intestinal permeability and visceral pain, along with certain behavioral features (Amini-Khoei et al., 2019; Labouesse et al., 2015; O'Mahony et al., 2011). Additionally, the animals exposed to early-life immune activation or psychological stress also exhibited intestinal dysbiosis (Amini-Khoei et al., 2019; Rincel et al., 2019).

Currently, there is a dearth of studies on schizophrenia and its associated microbiota. Our study examined the changes in specific types of bacteria, such as *Bacteroides* spp., *Bifidobacterium* spp., *Clostridium coccoides* group, *E. coli*, and *Lactobacillus* spp., in adult MIA rat offspring, when compared to control group. Our MIA rat model showed that some of these differences were consistent with the clinical case control study of schizophrenia. Higher incidences of lactic-acid bacteria in schizophrenia have been observed in the microbiota composition of oropharynx, when compared with controls (Castro-Nallar et al., 2015). Additionally, at the family level, *Lactobacillaceae* was increased in first-episode psychosis patients compared to controls (Schwarz et al., 2017).

A balanced intestinal environment with the microbiota plays an essential role in the health status of the host, by exerting beneficial effects on neurodevelopment and behavior phenotypes (Sharon et al., 2016). In a major subset of patients, intestinal symptoms arose before the onset of a mood disorder, suggesting that primary gut disturbance may serve as an underlying driver in some patient subgroups (Sharon et al., 2016). Another clue confirming the gut microbiota regulates behaviors comes from studies of germ-free mice (Chu et al., 2019). These mice tend to have hyperactivity disorder and risk-taking behaviors as well as memory and learning impairment when compared to conventional (specific-pathogen-free) mice (Clarke et al., 2013; Gareau et al., 2011; Diaz Heijt et al., 2011; Neufeld et al., 2011; Sharon et al., 2016). Previous findings have indicated that the gut microbiota can affect the expression levels of genes in the medial prefrontal cortex (mPFC) (Hoban et al., 2016, 2017). The mPFC can regulate specific emotional behaviors and mediate the effects of early-life stresses (Hao et al., 2019). Intestinal alterations, especially gut microbiota dysbiosis and loss of barrier functions, can affect brain development and permanently diminish gut-brain communications (Meyer, 2014).




FIGURE 5 Specific fecal bacteria measurements in MIA and control offspring. Total microbiota count (a) and the proportion of *E. coli* (b), *Lactobacillus* spp. (c), *Bifidobacterium* spp. (d), *Bacteroides* spp. (e), *Clostridium coccoides* group (f) and *Fusobacterium prausnitzii* (g) were analyzed and shown as the mean \pm SEM, $n = 6$ per group. Independent Student's *t*-test (two-tailed) was performed for statistical analysis. * $p < .05$, ** $p < .01$. The detection was repeated independently for three times

Long-range interactions between the brain and the gut microbiota have been reported (Bravo et al., 2011; Ochoa-Reparaz et al., 2010). A recent study reported GI barrier impairment and microbiota alteration in the MIA mice (Hsiao et al., 2013). Administration of *Bacteroides fragilis* to MIA offspring could improve gut permeability and microbial composition, as well as prevent abnormal communicative, stereotyped, sensorimotor, and anxiety-like behaviors (Hsiao et al., 2013). Numerous studies, using MIA or maternal separation models, have demonstrated that probiotic treatment and fecal transplantation can mediate brain function and behavioral features (Desbonnet et al., 2010; Hsiao et al., 2013). Some studies have suggested probiotics are effective in treating the emotional symptoms of chronic fatigue syndrome and psychological distress in humans (Messaoudi et al., 2011; Rao et al., 2009).

The potential roles of microbiota in the etiology of psychiatric disease should be tested. Growth studies have suggested that most of the effects of microbiota on brain are regulated by microbial metabolites. Kynurenine (Kyn) accumulation in the brain has been associated

with depression and schizophrenia (Muller & Schwarz, 2007); gut microbiota regulation of tryptophan (Trp) metabolism across three metabolic pathways can lead to the formation of serotonin, Kyn and indole derivatives (Cervenka et al., 2017). Although the blood-brain barrier is highly selective, Trp and Kyn derivatives can pass through this barrier and elicit noticeable effects on neurotransmitter function (Cervenka et al., 2017). Kynurenic acid and Quin, produced directly from Kyn, affect the brain differently by binding to glutamate receptors (e.g., N-methyl-D-aspartate receptor), which is essential for memory function (Cervenka et al., 2017). Short-chain fatty acids (SCFAs), which are important bacterial fermentation products, drive microglia maturation to fulfil the requirement for microglia maintenance (Erny et al., 2015). Interestingly, decreased SCFA levels in the feces of Parkinson's disease patients were recently reported (Unger et al., 2016). Previous research has suggested that treatment with 4-ethylphenylsulfate to naive wild-type mice can induce anxiety-like behaviors, as similar to those observed in MIA offspring (Hsiao et al., 2013).




FIGURE 6 The inflammatory response in MIA and control offspring. (a–c) Maternal Poly I:C treatment induced gut inflammatory responses in offspring as characterized by increased mRNA and protein expression of the pro-inflammatory cytokines TNF- α and IL-1 β in the colon as indicated by real-time PCR (a) and western blotting (b and c), with no significant changes was found in IL-6 expression (a–c). (d and e) Inflammatory cytokine responses in prefrontal cortex of MIA and control offspring. The protein expression of the pro-inflammatory cytokines IL-1 β , IL-6, and TNF- α in the prefrontal cortex of experimental animals as indicated by western blotting (d). Iba1 staining in the prefrontal cortex of MIA and control offspring (e). $n = 6$ for mRNA and protein expression and $n = 3$ for immunofluorescence staining for each group. Independent Student's *t*-test (two-tailed) was performed for statistical analysis. All data are expressed as the mean \pm SEM. * $p < .05$, ** $p < .01$. All experiments were repeated independently for three times

Neuroinflammation plays an essential role in the pathophysiology of neurodegenerative disease (Cappellano et al., 2013; Severance et al., 2013). Proinflammatory cytokines are secreted from the brain and peripheral neurons (McCoy & Tansey, 2008). The passage of microbially associated molecular patterns from the intestine to the brain can result in low inflammation levels (Pal et al., 2015). Such persistent proinflammatory signaling is associated with the severity of neurodegenerative disease (Cappellano et al., 2013). However, our results indicated that no obvious changes in the levels of proinflammatory cytokines were found in the prefrontal cortex of MIA offspring, although increased proinflammatory cytokine levels were detected in the intestines. Microglia are the major sources of cytokine and chemokine expression in the brain parenchyma; however, our results showed no obvious microglial anomalies in the prefrontal cortex of MIA offspring. Recent work has reported that multiple microglial and astrocyte markers remained unchanged in the midbrains of MIA offspring, as detected by Western blotting, even though their mRNA levels were increased (Purves-Tyson et al., 2019). These authors explained that RNA detection may be more sensitive for immune activation (Purves-Tyson et al., 2019), which could also support our divergent results. Another study indicated Poly I:C significantly increased microglial activation at PND21 in male hippocampi,

which support the hypothesis of neuroinflammation of MIA (Murray et al., 2019). The brain inflammatory state of MIA offspring may vary in different brain regions or stage of development.

This study was limited as it was an observational study, and the association between intestinal changes and behavioral deficits was not determined. It is unclear whether gut microbiota can confer risk to neuropsychiatric disorders or is a consequence of pathological processes. Therefore, future investigation is needed to understand the intricate and precise interactions between a host and its associated microbial communities. Another limitation of this study is that only male rats were analyzed and future research need to avoid sex bias.

ACKNOWLEDGMENTS

LXL and WQL designed the study and wrote the protocol. XS established the animal model of MIA and wrote the manuscript. MXC helped in sample preparation and interpretation of the results. XF, MS, MLS, YFY, LWZ and QL undertook the statistical analysis and helped with molecular biology techniques. All authors contributed to and have approved the final manuscript. This work was supported by the National Natural Science Foundation of China (82001407 to XS; 81972152 to LL; U1904130 WL.); Medical

Science and Technology Research Project of Henan Province (2018020371 to XS; LHGJ20190476 to MS; LHGJ20190482 to QL); Key Scientific Research Project of Henan Higher Education Institutions (20B320016 to MS); High Scientific and Technological Research Fund of Xinxiang Medical University (2017ZDCG-04 to LL); The Open Fund of Henan Key lab of Biological Psychiatry (ZDSYS2018008 to XS; ZDSYS2018003 to MS).

CONFLICT OF INTEREST

The authors declare that they have no competing interests.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1002/brb3.2133>.

DATA AVAILABILITY STATEMENT

The datasets used and/or analyzed during the current study are available from the corresponding authors upon reasonable request.

ORCID

Wenqiang Li  <https://orcid.org/0000-0002-0833-246X>

Mengxue Chen  <https://orcid.org/0000-0003-1751-5759>

REFERENCES

- Akhondzadeh, S. (2019). Microbiome and schizophrenia. *Avicenna Journal of Medical Biotechnology*, 11(4), 269.
- Amini-Khoei, H., Haghani-Samani, E., Beigi, M., Soltani, A., Mobini, G. R., Balali-Dehkordi, S., Haj-Mirzaian, A., Rafieian-Kopaei, M., Alizadeh, A., Hojjati, M. R., & Validi, M. (2019). On the role of corticosterone in behavioral disorders, microbiota composition alteration and neuroimmune response in adult male mice subjected to maternal separation stress. *International Immunopharmacology*, 66, 242–250. <https://doi.org/10.1016/j.intimp.2018.11.037>
- Bergdolt, L., & Dunaevsky, A. (2019). Brain changes in a maternal immune activation model of neurodevelopmental brain disorders. *Progress in Neurobiology*, 175, 1–19. <https://doi.org/10.1016/j.pneurobio.2018.12.002>
- Bravo, J. A., Forsythe, P., Chew, M. V., Escaravage, E., Savignac, H. M., Dinan, T. G., Bienenstock, J., & Cryan, J. F. (2011). Ingestion of Lactobacillus strain regulates emotional behavior and central GABA receptor expression in a mouse via the vagus nerve. *Proceedings of the National Academy of Sciences*, 108(38), 16050–16055. <https://doi.org/10.1073/pnas.1102999108>
- Brown, A. S., Begg, M. D., Gravenstein, S., Schaefer, C. A., Wyatt, R. J., Bresnahan, M., Babulas, V. P., & Susser, E. S. (2004). Serologic evidence of prenatal influenza in the etiology of schizophrenia. *Archives of General Psychiatry*, 61(8), 774–780. <https://doi.org/10.1001/archpsyc.61.8.774>
- Brown, A. S., & Meyer, U. (2018). Maternal immune activation and neuropsychiatric illness: A translational research perspective. *American Journal of Psychiatry*, 175(11), 1073–1083. <https://doi.org/10.1176/appi.ajp.2018.17121311>
- Cappellano, G., Carecchio, M., Fleetwood, T., Magistrelli, L., Cantello, R., Dianzani, U., & Comi, C. (2013). Immunity and inflammation in neurodegenerative diseases. *American Journal of Neurodegenerative Disease*, 2(2), 89–107.
- Castro-Nallar, E., Bendall, M. L., Pérez-Losada, M., Sabuncyan, S., Severance, E. G., Dickerson, F. B., Schroeder, J. R., Yolken, R. H., & Crandall, K. A. (2015). Composition, taxonomy and functional diversity of the oropharynx microbiome in individuals with schizophrenia and controls. *PeerJ*, 3, e1140. <https://doi.org/10.7717/peerj.1140>
- Cervenka, I., Agudelo, L. Z., & Ruas, J. L. (2017). Kynurenines: Tryptophan's metabolites in exercise, inflammation, and mental health. *Science*, 357(6349), eaaf9794. <https://doi.org/10.1126/science.aaf9794>
- Chu, C., Murdock, M. H., Jing, D., Won, T. H., Chung, H., Kressel, A. M., Tsaava, T., Addorisio, M. E., Putzel, G. G., Zhou, L., Bessman, N. J., Yang, R., Moriyama, S., Parkhurst, C. N., Li, A., Meyer, H. C., Teng, F., Chavan, S. S., Tracey, K. J., ... Artis, D. (2019). The microbiota regulate neuronal function and fear extinction learning. *Nature*, 574(7779), 543–548. <https://doi.org/10.1038/s41586-019-1644-y>
- Clarke, G., Grenham, S., Scully, P., Fitzgerald, P., Moloney, R. D., Shanahan, F., Dinan, T. G., & Cryan, J. F. (2013). The microbiome-gut-brain axis during early life regulates the hippocampal serotonergic system in a sex-dependent manner. *Molecular Psychiatry*, 18(6), 666–673. <https://doi.org/10.1038/mp.2012.77>
- Cryan, J. F., & Dinan, T. G. (2012). Mind-altering microorganisms: The impact of the gut microbiota on brain and behaviour. *Nature Reviews Neuroscience*, 13(10), 701–712. <https://doi.org/10.1038/nrn3346>
- Desbonnet, L., Garrett, L., Clarke, G., Kiely, B., Cryan, J. F., & Dinan, T. G. (2010). Effects of the probiotic Bifidobacterium infantis in the maternal separation model of depression. *Neuroscience*, 170(4), 1179–1188. <https://doi.org/10.1016/j.neuroscience.2010.08.005>
- Erny, D., Hrabě de Angelis, A. L., Jaitin, D., Wieghofer, P., Staszewski, O., David, E., Keren-Shaul, H., Mhlahkoi, T., Jakobshagen, K., Buch, T., Schwierzeck, V., Utermöhlen, O., Chun, E., Garrett, W. S., McCoy, K. D., Diefenbach, A., Staeheli, P., Stecher, B., Amit, I., & Prinz, M. (2015). Host microbiota constantly control maturation and function of microglia in the CNS. *Nature Neuroscience*, 18(7), 965–977. <https://doi.org/10.1038/nn.4030>
- Gareau, M. G., Wine, E., Rodrigues, D. M., Cho, J. H., Whary, M. T., Philpott, D. J., MacQueen, G., & Sherman, P. M. (2011). Bacterial infection causes stress-induced memory dysfunction in mice. *Gut*, 60(3), 307–317. <https://doi.org/10.1136/gut.2009.202515>
- Golofast, B., & Vales, K. (2020). The connection between microbiome and schizophrenia. *Neuroscience and Biobehavioral Reviews*, 108, 712–731. <https://doi.org/10.1016/j.neubiorev.2019.12.011>
- Hao, K., Su, X. I., Luo, B., Cai, Y., Chen, T., Yang, Y., Shao, M., Song, M., Zhang, L., Zhong, Z., Li, W., & Lv, L. (2019). Prenatal immune activation induces age-related alterations in rat offspring: Effects upon NMDA receptors and behaviors. *Behavioral Brain Research*, 370, 111946. <https://doi.org/10.1016/j.bbr.2019.111946>
- Heijtz, R. D., Wang, S., Anuar, F., Qian, Y., Bjorkholm, B., Samuelsson, A., Hibberd, M. L., Forssberg, H., & Pettersson, S. (2011). Normal gut microbiota modulates brain development and behavior. *Proceedings of the National Academy of Sciences*, 108(7), 3047–3052. <https://doi.org/10.1073/pnas.1010529108>
- Hoban, A. E., Stilling, R. M., M. Moloney, G., Moloney, R. D., Shanahan, F., Dinan, T. G., Cryan, J. F., & Clarke, G. (2017). Microbial regulation of microRNA expression in the amygdala and prefrontal cortex. *Microbiome*, 5(1), 102. <https://doi.org/10.1186/s40168-017-0321-3>
- Hoban, A. E., Stilling, R. M., Ryan, F. J., Shanahan, F., Dinan, T. G., Claesson, M. J., Clarke, G., & Cryan, J. F. (2016). Regulation of prefrontal cortex myelination by the microbiota. *Translational Psychiatry*, 6, e774. <https://doi.org/10.1038/tp.2016.42>
- Hsiao, E. Y., McBride, S. W., Hsien, S., Sharon, G., Hyde, E. R., McCue, T., Codelli, J. A., Chow, J., Reisman, S. E., Petrosino, J. F., Patterson, P. H., & Mazmanian, S. K. (2013). Microbiota modulate behavioral and physiological abnormalities associated with neurodevelopmental disorders. *Cell*, 155(7), 1451–1463. <https://doi.org/10.1016/j.cell.2013.11.024>

- Hui, C. W., St-Pierre, A., El Hajj, H., Remy, Y., Hébert, S. S., Luheshi, G. N., Srivastava, L. K., & Tremblay, M.-È. (2018). Prenatal immune challenge in mice leads to partly sex-dependent behavioral, microglial, and molecular abnormalities associated with schizophrenia. *Frontiers in Molecular Neuroscience*, *11*, 13. <https://doi.org/10.3389/fnmol.2018.00013>
- Labouesse, M. A., Langhans, W., & Meyer, U. (2015). Long-term pathological consequences of prenatal infection: Beyond brain disorders. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology*, *309*(1), R1–R12. <https://doi.org/10.1152/ajpregu.00087.2015>
- Mayer, E. A. (2011). Gut feelings: The emerging biology of gut-brain communication. *Nature Reviews Neuroscience*, *12*(8), 453–466. <https://doi.org/10.1038/nrn3071>
- Mayer, E. A., Knight, R., Mazmanian, S. K., Cryan, J. F., & Tillisch, K. (2014). Gut microbes and the brain: Paradigm shift in neuroscience. *Journal of Neuroscience*, *34*(46), 15490–15496. <https://doi.org/10.1523/JNEUROSCI.3299-14.2014>
- McCoy, M. K., & Tansey, M. G. (2008). TNF signaling inhibition in the CNS: Implications for normal brain function and neurodegenerative disease. *Journal of Neuroinflammation*, *5*, 45. <https://doi.org/10.1186/1742-2094-5-45>
- Mednick, S. A., Machon, R. A., Huttunen, M. O., & Bonett, D. (1988). Adult schizophrenia following prenatal exposure to an influenza epidemic. *Archives of General Psychiatry*, *45*(2), 189–192. <https://doi.org/10.1001/archpsyc.1988.01800260109013>
- Meehan, C., Harms, L., Frost, J. D., Barreto, R., Todd, J., Schall, U., Shannon Weickert, C., Zavitsanou, K., Michie, P. T., & Hodgson, D. M. (2017). Effects of immune activation during early or late gestation on schizophrenia-related behaviour in adult rat offspring. *Brain, Behavior, and Immunity*, *63*, 8–20. <https://doi.org/10.1016/j.bbi.2016.07.144>
- Messaoudi, M., Lalonde, R., Violle, N., Javelot, H., Desor, D., Nejd, A., & Cazaubiel, J. M. (2011). Assessment of psychotropic-like properties of a probiotic formulation (*Lactobacillus helveticus* R0052 and *Bifidobacterium longum* R0175) in rats and human subjects. *British Journal of Nutrition*, *105*(5), 755–764. <https://doi.org/10.1017/S0007114510004319>
- Meyer, U. (2014). Prenatal poly(i:C) exposure and other developmental immune activation models in rodent systems. *Biological Psychiatry*, *75*(4), 307–315. <https://doi.org/10.1016/j.biopsych.2013.07.011>
- Meyer, U., & Feldon, J. (2010). Epidemiology-driven neurodevelopmental animal models of schizophrenia. *Progress in Neurobiology*, *90*(3), 285–326. <https://doi.org/10.1016/j.pneurobio.2009.10.018>
- Meyer, U., Feldon, J., Schedlowski, M., & Yee, B. K. (2006). Immunological stress at the maternal-foetal interface: A link between neurodevelopment and adult psychopathology. *Brain, Behavior, and Immunity*, *20*(4), 378–388. <https://doi.org/10.1016/j.bbi.2005.11.003>
- Meyer, U., Nyffeler, M., Engler, A., Urwyler, A., Schedlowski, M., Knuesel, I., & Feldon, J. (2006). The time of prenatal immune challenge determines the specificity of inflammation-mediated brain and behavioral pathology. *Journal of Neuroscience*, *26*(18), 4752–4762. <https://doi.org/10.1523/JNEUROSCI.0099-06.2006>
- Meyer, U., Nyffeler, M., Schwendener, S., Knuesel, I., Yee, B. K., & Feldon, J. (2008). Relative prenatal and postnatal maternal contributions to schizophrenia-related neurochemical dysfunction after in utero immune challenge. *Neuropsychopharmacology*, *33*(2), 441–456. <https://doi.org/10.1038/sj.npp.1301413>
- Minakova, E., & Warner, B. B. (2018). Maternal immune activation, central nervous system development and behavioral phenotypes. *Birth Defects Research*, *110*(20), 1539–1550. <https://doi.org/10.1002/bdr2.1416>
- Muller, N., & Schwarz, M. J. (2007). The immune-mediated alteration of serotonin and glutamate: Towards an integrated view of depression. *Molecular Psychiatry*, *12*(11), 988–1000. <https://doi.org/10.1038/sj.mp.4002006>
- Murray, K. N., Edey, M. E., Manca, M., Vernon, A. C., Oladipo, J. M., Fasolino, V., Harte, M. K., Mason, V., Grayson, B., McHugh, P. C., Knuesel, I., Prinssen, E. P., Hager, R., & Neill, J. C. (2019). Evolution of a maternal immune activation (mIA) model in rats: Early developmental effects. *Brain, Behavior, and Immunity*, *75*, 48–59. <https://doi.org/10.1016/j.bbi.2018.09.005>
- Nemani, K., Hosseini Ghomi, R., McCormick, B., & Fan, X. (2015). Schizophrenia and the gut-brain axis. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *56*, 155–160. <https://doi.org/10.1016/j.pnpbp.2014.08.018>
- Neufeld, K. M., Kang, N., Bienenstock, J., & Foster, J. A. (2011). Reduced anxiety-like behavior and central neurochemical change in germ-free mice. *Neurogastroenterology and Motility*, *23*(3), 255–264. <https://doi.org/10.1111/j.1365-2982.2010.01620.x>
- Nguyen, T. T., Kosciolk, T., Maldonado, Y., Daly, R. E., Martin, A. S., McDonald, D., Knight, R., & Jeste, D. V. (2019). Differences in gut microbiome composition between persons with chronic schizophrenia and healthy comparison subjects. *Schizophrenia Research*, *204*, 23–29. <https://doi.org/10.1016/j.schres.2018.09.014>
- Ochoa-Reparaz, J., Mielcarz, D. W., Ditrio, L. E., Burroughs, A. R., Begum-Haque, S., Dasgupta, S., & Kasper, L. H. (2010). Central nervous system demyelinating disease protection by the human commensal *Bacteroides fragilis* depends on polysaccharide A expression. *The Journal of Immunology*, *185*(7), 4101–4108. <https://doi.org/10.4049/jimmunol.1001443>
- O'Mahony, S. M., Hyland, N. P., Dinan, T. G., & Cryan, J. F. (2011). Maternal separation as a model of brain-gut axis dysfunction. *Psychopharmacology (Berl)*, *214*(1), 71–88. <https://doi.org/10.1007/s00213-010-2010-9>
- Pal, G. D., Shaikh, M., Forsyth, C. B., Ouyang, B., Keshavarzian, A., & Shannon, K. M. (2015). Abnormal lipopolysaccharide binding protein as marker of gastrointestinal inflammation in Parkinson disease. *Frontiers in Neuroscience*, *9*, 306. <https://doi.org/10.3389/fnins.2015.00306>
- Purves-Tyson, T. D., Weber-Stadlbauer, U., Richetto, J., Rothmond, D. A., Labouesse, M. A., Polesel, M., Robinson, K., Shannon Weickert, C., & Meyer, U. (2019). Increased levels of midbrain immune-related transcripts in schizophrenia and in murine offspring after maternal immune activation. *Molecular Psychiatry*, *26*(3), 849–863. <https://doi.org/10.1038/s41380-019-0434-0>
- Rahman, T., Zavitsanou, K., Purves-Tyson, T., Harms, L. R., Meehan, C., Schall, U., Todd, J., Hodgson, D. M., Michie, P. T., & Weickert, C. S. (2017). Effects of immune activation during early or late gestation on N-Methyl-D-aspartate receptor measures in adult rat offspring. *Frontiers in Psychiatry*, *8*, 77. <https://doi.org/10.3389/fpsy.2017.00077>
- Rao, A. V., Bsted, A. C., Beaulne, T. M., Katzman, M. A., Iorio, C., Berardi, J. M., & Logan, A. C. (2009). A randomized, double-blind, placebo-controlled pilot study of a probiotic in emotional symptoms of chronic fatigue syndrome. *Gut Pathogens*, *1*(1), 6. <https://doi.org/10.1186/1757-4749-1-6>
- Rincel, M., Aubert, P., Chevalier, J., Grohard, P.-A., Basso, L., Monchaux de Oliveira, C., Helbling, J. C., Lévy, É., Chevalier, G., Leboyer, M., Eberl, G., Layé, S., Capuron, L., Vergnolle, N., Neunlist, M., Boudin, H., Lepage, P., & Darnaudéry, M. (2019). Multi-hit early life adversity affects gut microbiota, brain and behavior in a sex-dependent manner. *Brain, Behavior, and Immunity*, *80*, 179–192. <https://doi.org/10.1016/j.bbi.2019.03.006>
- Schwarz, E., Maukonen, J., Hyytiäinen, T., Kieseppä, T., Orešič, M., Sabunciyani, S., Mantere, O., Saarela, M., Yolken, R., & Suvisaari, J. (2017). Analysis of microbiota in first episode psychosis identifies preliminary associations with symptom severity and treatment response. *Schizophrenia Research*, *192*, 398–403.

- Severance, E. G., Gressitt, K. L., Stallings, C. R., Origoni, A. E., Khushalani, S., Leweke, F. M., Dickerson, F. B., & Yolken, R. H. (2013). Discordant patterns of bacterial translocation markers and implications for innate immune imbalances in schizophrenia. *Schizophrenia Research*, 148(1-3), 130-137. <https://doi.org/10.1016/j.schres.2013.05.018>
- Sharon, G., Sampson, T. R., Geschwind, D. H., & Mazmanian, S. K. (2016). The central nervous system and the gut microbiome. *Cell*, 167(4), 915-932. <https://doi.org/10.1016/j.cell.2016.10.027>
- Stefano, G. B., Pilonis, N., Ptacek, R., Raboch, J., Vnukova, M., & Kream, R. M. (2018). Gut, microbiome, and brain regulatory axis: Relevance to neurodegenerative and psychiatric disorders. *Cellular and Molecular Neurobiology*, 38(6), 1197-1206. <https://doi.org/10.1007/s10571-018-0589-2>
- Tillisch, K., Labus, J., Kilpatrick, L., Jiang, Z., Stains, J., Ebrat, B., Guyonnet, D., Legrain-Raspaud, S., Trotin, B., Naliboff, B., & Mayer, E. A. (2013). Consumption of fermented milk product with probiotic modulates brain activity. *Gastroenterology*, 144(7), 1394-1401, 1401 e1391-1394.
- Unger, M. M., Spiegel, J., Dillmann, K.-U., Grundmann, D., Philippeit, H., Bürmann, J., Faßbender, K., Schwiertz, A., & Schäfer, K.-H. (2016). Short chain fatty acids and gut microbiota differ between patients with Parkinson's disease and age-matched controls. *Parkinsonism & Related Disorders*, 32, 66-72. <https://doi.org/10.1016/j.parkrel.2016.08.019>
- Van Oudenhove, L., Vandenberghe, J., Demyttenaere, K., & Tack, J. (2010). Psychosocial factors, psychiatric illness and functional gastrointestinal disorders: A historical perspective. *Digestion*, 82(4), 201-210. <https://doi.org/10.1159/000269822>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Li W, Chen M, Feng X, et al. Maternal immune activation alters adult behavior, intestinal integrity, gut microbiota and the gut inflammation. *Brain Behav.* 2021;11:e02133. <https://doi.org/10.1002/brb3.2133>